



-Publisher ; W. Junk, [1979]

Bass (/eds?search=y&query=%22Bass%22&type=MM)/\*parasitology  
 (/eds?search=y&query=%22Bass+parasitology%22&type=MM) ;  
 Copepoda (/eds?search=y&query=%22Copepoda%22&type=MM)  
     /\*classification (/eds?search=y&  
     query=%22Copepoda+classification%22&type=MM); Animals  
     (/eds?search=y&query=%22Animals%22&type=MH) ; Copepoda  
     (/eds?search=y&query=%22Copepoda%22&type=MH)/anatomy &  
     histology (/eds?search=y&  
     query=%22Copepoda+anatomy+%26+histology%22&type=MH) ; Female  
     (/eds?search=y&query=%22Female%22&type=MH) ; Gills  
     (/eds?search=y&query=%22Gills%22&type=MH)/parasitology  
     (/eds?search=y&query=%22Gills+parasitology%22&type=MH) ;  
     Mediterranean Sea (/eds?search=y&  
     query=%22Mediterranean+Sea%22&type=MH) ; Species Specificity  
     (/eds?search=y&query=%22Species+Specificity%22&type=MH)

### **:Subject Terms**

A new species of parasitic copepod, *Caligus tunisiensis* n. sp. (Caligidae), is described based on two female specimens collected from the gills of the painted comber, *Serranus scriba* (L.), caught in the Mediterranean Sea, off the Tunisian coasts. The new species belongs to the *Caligus productus*-species group established by Boxshall & Gurney (1980) as it shares the following set of character states: (i) antenna with well-developed posterior process on proximal segment; (ii) posterior margin of distal exopodal segment of leg 1 lacking typical plumose setae, or retaining single vestigial seta; and (iii) 2-segmented exopod of leg 4 armed with IV spines on compound distal exopodal segment. Detailed morphological comparisons between the new species and the core members of the *C. productus*-species group revealed that the new species closely resembles with *C. productus* Dana, 1852 and *C. temnodontis* Brian, 1924. However, the new species can be distinguished from its congeners in having: (i) a female maxilliped bearing a prominent bi-lobate myxal process, opposing tip of the claw; (ii) leg 4 protopod ornamented with a patch of spinules on the posterolateral surface; and (iii) an abdomen ornamented with two rows of minute spinules at the posterolateral corners.

Bahri, L. (1995). Analyse génétique et parasitologique des populations du loup commun, *Dicentrarchus labrax* et du loup ponctué *Dicentrarchus punctatus* des côtes tunisiennes. MSc thesis, Université de Tunis El Manar, 121 pp

### **:References**

Ben Hassine, O. K. (1983). Les Copépodes parasites de poissons Mugilidae en Méditerranée occidentale (côtes françaises et tunisiennes).

Morphologie, Bio-écologie, cycles évolutifs. PhD thesis, U.S.T.L.,  
Montpellier, 452 pp

Benmansour, B. (2001). Biodiversité et bioécologie des copépodes parasites des poissons téléostéens. PhD thesis, Université de Tunis El Manar, 454 pp

Benmansour, B., & Ben Hassine, O. K. (1997). Première mention en Tunisie de certains Caligidae et Lernaeopodidae (Copepoda) parasites de poissons Téléostéens. *Acta Ichthyologica et Piscatoria*, 20, 157–175  
Benmansour, B., & Ben Hassine, O. K. (1998). Preliminary analysis of parasitic copepod species richness among coastal fishes of Tunisia.  
:Italian Journal of Zoology, 65, 341–344. (PMID  
(10.1080/11250009809386844

Boxshall, G. A. (1990). The skeletomusculature of siphonostomatoid copepods, with an analysis of adaptive radiation in structure of the oral cone. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 328, 167–212

Boxshall, G. A. (2018). The sea lice (Copepoda: Caligidae) of Moreton Bay (Queensland, Australia) with descriptions of thirteen new species.  
(zootaxa.4398.1.1/10.11646 :Zootaxa, 4398, 1–172. (PMID

Boxshall, G. A., & El-Rashidy, H. H. (2009). A review of the *Caligus productus* species group, with the description of a new species, new synonymies and supplementary descriptions. *Zootaxa*, 2271, 1–26.  
(zootaxa.2271.1.1/10.11646 :(PMID

Boxshall, G. A., & Gurney, A. R. (1980). Description of two new and one poorly known species of the genus *Caligus* Müller, 1785 (Copepoda: Siphonostomatoida). *Bulletin of the British Museum (Natural History)*, 39,  
.161–178

Cabral, P., & Raibaut, A. (1986). Découverte d'un copépode caligide nouveau, parasite du tégument du loup, *Dicentrarchus labrax* (L., 1758)  
(Pisces, Moronidae) en élevage et en milieu naturel. *Bulletin de la Société Zoologique de France*, 111, 123–130

Essafi, K., Cabral, P., & Raibaut, A. (1984). Copépodes parasites de poissons des îles Kerkennah (Tunisie méridionale). Les archives de l'Institut Pasteur Tunis, 61, 475–523

Froese, R., & Pauly, D. (2020). FishBase. World Wide Web electronic publication. [www.fishbase.org](http://www.fishbase.org) . Accessed on 14 April 2020

Huys, R., & Boxshall, G. A. (1991). Copepod Evolution. London: The Ray Society

ICZN (2012). International Commission on Zoological Nomenclature: Amendment of articles 8, 9, 10, 21 and 78 of the International Code of Zoological Nomenclature to expand and refine methods of publication.  
:Bulletin of Zoological Nomenclature, 69, 161–169. (PMID  
(bzn.v69i3.a8.161/10.21805

- Kamanli, S. A., Kihara, T. C., Ball, A. D., Morritt, D., & Clark, P. F. (2017). A 3D imaging and visualisation workflow, using confocal microscopy and advanced image processing for brachyuran crab larvae. *Journal of Microscopy*, 266, 307–323. (PMID 283211288)
- Limaye, A. (2012). Drishti: a volume exploration and presentation tool. In: Stock, S. R. (Ed.) *Proceedings of SPIE Vol. 8506. Developments in X-Ray Tomography VIII*, 85060X, 17 October 2012, San Diego, California, USA
- Ohtsuka, S., & Boxshall, G. A. (2019). Two new species of the genus *Caligus* (Crustacea, Copepoda, Siphonostomatoida) from the Sea of Japan, with a note on the establishment of a new species group. (*Zookeys*.893.46923/10.3897 :ZooKeys, 893, 91–113. (PMID 31189288))
- Özak, A. A., Sakarya, Y., & Boxshall, G. A. (2019). *Caligus adanensis* n. sp. (Copepoda: Caligidae Burmeister, 1835) parasitic on garfish, *Belone belone* (Linnaeus, 1760), from the eastern Mediterranean Sea, off the Turkish coast. *Marine Biodiversity*, 49, 1877–1890. (PMID 312526-019-00949-9/10.1007)
- Papoutsoglou, S. E. (1976). Metazoan parasites of fishes from Saronicos Gulf, Athens, Greece. *Thalassographica*, 1, 69–102
- Raibaut, A., Ben Hassine, O. K., & Maamouri, K. (1971). Copépodes parasites des poissons de Tunisie (première série). *Bulletin de l'Institut National Scientifique et Technique d'Océanographie et de Pêche*, 2, 169–197
- Raibaut, A., & Ben Hassine, O. K. (1977). Les Copépodes parasites des Muges en Méditerranée. *Bulletin du Muséum National d'Histoire Naturelle*, 329, 833–848
- Rodrigues, A. M. V., Özak, A. A., Silva, L. M. H., & Boxshall, G. A. (2018). *Caligus mulli* n. sp. (Copepoda: Caligidae) parasitic on two Mullid fishes from the eastern Mediterranean and adjacent Atlantic waters. (*Parasitology Research*, 117, 3843–3850. (PMID 300436-018-6090-5/10.1007))
- Walter, T. C., & Boxshall, G. A. (2020). World of Copepods Database. <http://www.marinespecies.org/copepoda> on 2020-04-06. Accessed on 10 April 2020
- Yamak, S., & Ben Hassine, O. K. (2002). Impact of a fish farm on ichthyofauna and ichthyoparasitofauna of a Tunisian lagoon. In: *Proceedings of the International Conference on the Environmental Problems of the Mediterranean Region*, 12–15 April 2002, Nicosia, Turkish Republic of Northern Cyprus
- Youssef, F., Benmansour, B., Ben Hassine, O. K., & Zouari-Tlig, S. (2016). Some parasitic copepods of selected teleost and chondrichthyan fishes from the Tunisian Gulfs. *African Journal of Microbiology Research*, (AJMR2015.7820/10.5897 :10, 1467–1476. (PMID 27342888))

*Date Created:* 20201119 *Date Completed:* 20210909 *Latest Revision:* 20210909 **:Publication Date**

20211105 **:Publication Date**

s11230-020-09959-9/10.1007 **:Accession Number**

33211288 **:Accession Number**