

Check Storage



Borrower:

SCIENCE SHELVED BY TITLE n.s.v.3 1969-70 (LIMITED LOAN)
MISSINGSCIENCE
SHELVED BY TITLE n.s.v.4 1971-72 (LIMITED LOAN) MISSING
SCIENCE SHELVED BY TITLE n.s.v.5-6 1973-75 (LIMITED LOAN)
MISSINGSCIENCE
SHELVED BY TITLE n.s.v.7-8 1976 (LIMITED LOAN) MISSING
SCIENCE SHELVED BY TITLE n.s.v.9 1977 (inc) (LIMITED LOAN)
MISSINGSCIENCE

1

131.94.29.30/131.94.29.29



Record 24 of 101
ILL pe

Record 44 of 101

CAN YOU SUPPLY ? YES NO COND FUTUREDATE
:ILL: 3596000 :Borrower: FXG :ReqDate: 20040225 :NeedBefore: 20040326
:Status: IN PROCESS 20040302 :RecDate: :RenewalReq:
:OCLC: 1629885 :Source: OCLCILL :DueDate: N/A :NewDueDate:
:Lender: ILU,*ORU,VPI,NDD,IXA
:CALLNO: *Lender's OCLC LDR: ~~3-22-1969-1987~~
:TITLE: Monitore zoologico italiano. *Supplemento.*
:IMPRINT: Firenze : Valsalva Editrice, 1966-1987.
:ARTICLE: Dumont H.J.: On a collection of zooplankton from Somalia with
description of three new species of Copepoda
:VOL: 14 :NO: 7 :DATE: 1981 :PAGES: 103-111
:VERIFIED: <TN:97176>OCLC ISSN: 0374-9444 [Format: Serial]
:PATRON: Bruno, Maria Cristina
:SHIP TO:

N/A

Storage

\$ 8IFM

Interlibrary Loan, GL-273 Florida International University
Tamiami Trail
Miami, FL 33199 USA

*Ariel
3-2
82*

:BILL TO: (see above) Your Fein# REQUIRED on bills. FXG's FEIN#59-6001874
:SHIP VIA: Library Rate or DLLI (FLIN only) :MAXCOST: \$30IFM
:COPYRT COMPLIANCE: CCL
:FAX: (305)348-6055, ARIEL 131.94.29.30 or 131.94.29.29
:E-MAIL: upill@fiu.edu
:BILLING NOTES: ILL Order numbers REQUIRED on invoices for processing!
:BORROWING NOTES: Attn ARIEL Libraries -- receiving docs via Ariel as a
borrower? Then pls send docs via Ariel as a lender. Thanks!
:AFFILIATION: SEFLIN,FLIN/DLLI,SOLINE/SO6,ASERL,LVIS,various reciprocals

OCLC Number:

Interlibrary Loan, GL-273 Florida International University
Tamiami Trail
Miami, FL 33199 USA

Monitore zoologico italiano

ITALIAN JOURNAL OF ZOOLOGY

PUBBLICATO DALLA UNIVERSITÀ DEGLI STUDI DI FIRENZE
CON IL CONTRIBUTO DEL CONSIGLIO NAZIONALE DELLE RICERCHE

N. S. SUPPLEMENTO XIV

15.5.1981

NO. 7 : 103-111

ON A COLLECTION OF ZOOPLANKTON FROM SOMALIA, WITH A DESCRIPTION OF THREE NEW SPECIES OF COPEPODA

(PUBBLICAZIONI DEL CENTRO DI STUDIO
PER LA FAUNISTICA ED ECOLOGIA TROPICALI DEL C.N.R.: CXCIV)

HENRI J. DUMONT

Zoological Institute, University of Gent

Received 15 April 1980

Since no information is available on the zooplankton of the Horn of Africa, a small set of freshwater samples, collected and kindly made available by the Centro di Studio per la Faunistica ed Ecologia Tropicali of the Consiglio Nazionale delle Ricerche, Florence (Director, Prof. L. Pardi), provided an opportunity to put on record the first species of Cladocera and Copepoda from that part of Africa.

The scarcity of surface waters in Somalia by no means implies that the local planktonic fauna is uninteresting. The fact that three out of four copepod species found here are new to science, points towards a long history of uninterrupted and undisturbed evolution. On the other hand, the fact that at least one species was also found in samples from the Kenyan rift valley, suggests that past climatic phases of increased humidity permitted aquatic forms to disperse over relatively wide areas that are now at least partly arid or even hyperarid.

SUBORDER CLADOCERA

Moina brachiata (Jurine, 1850) (Fig. 1).

Material examined. — Maas (04°23'N-46°05'E), well near to the village, several ♀♀ MF (1) 008, leg. SBS (2), 1.IX.1968.

(1) MF = Museo Zoologico dell'Università di Firenze.

(2) SBS = Spedizione Biologica in Somalia del Centro di Studio per la Faunistica ed Ecologia Tropicali del C.N.R. (see PARDI, 1976).




Fig. 1. — Distribution of *Moina brachiata* (Jurine, 1850) in Africa and around the Mediterranean basin. Somalian record indicated by a square.

Remarks. — This species is rather widespread in Europe, North Africa including the central Sahara, and South Africa. Records are presented in Fig. 1 (from DUMONT, 1979). In eastern Africa, there was only one old and somewhat doubtful record from Lake Victoria (DADAY, 1907), that now

gains considerable credibility in the same general area. Most records for the pecten on the end of the antenna are of the unusually low number

Bosmina longirostris O.F.S.

Material examined. — leg. SBS, 1.IX.1968.

Remarks. — A single record amidst the individuals of


SUBC

Nitocrella somalica n. sp.

Material examined. — holotype (a fully dissected female), 7 ♀♀ MF 011

Description. — Total length 1.5 mm. Head of eight segments, bearing 1-4 all three-segmented. P₁-P₃, five appendages on each, with one inner spine on P₁ and with an inner spine on P₂ and P₃. P₄ segmented, with their basal strong spine on P₂-P₄. In a female, the spine, is seen. P₅: strong, blunt spines. Exopodites prominent over the basoendites. Spines, subequal in length, tapering towards their tip, and the basal spines stand at their vertex and bears five strong spines

Relationship. — Geographically from the beaches of Lake Victoria. In revision, it belongs to the



ine, 1850) in Africa and around the
d indicated by a square.

r widespread in Europe, North
uth Africa. Records are presented
n Africa, there was only one old
Victoria (DADAY, 1907), that now

gains considerable credibility by the discovery of a second population in the same general area. Morphologically, Somalian animals are noteworthy for the pecten on the end-claws of the postabdomen, which is composed of the unusually low number of only eight teeth.

Bosmina longirostris O.F. Müller, 1846.

Material examined. — Maas, well near to the village, 1 ♀ MF 009, leg. SBS, 1.IX.1968.

Remarks. — A single, somewhat damaged specimen was found amidst the individuals of the foregoing species.

Subclass COPEPODA

SUBORDER HARPACTICOIDA


Nitocrella somalica n. sp. (Figs 2-10).

Material examined. — Baidoa (03°07'N-43°38'E), well of the aqueduct; holotype (a fully dissected female, mounted in glycerin), 1 ♀ MF 010; paratypes, 7 ♀ ♀ MF 011; leg. SBS, 27.VIII.1968.

Description. — Total length around 0.5 mm. First antenna composed of eight segments, bearing a long esthete on top of segment 4. Exopodite of second antenna bearing three appendages. Exopodites of thoracopods 1-4 all three-segmented. Their apical segment with four appendages on P₁-P₃, five appendages on P₄. Middle segment without inner spine on P₁, with one inner spine on P₂-P₄. The endopodite of P₁ three-segmented and with an inner spine on the basal segment. Endopodites of P₂-P₄ two-segmented, with their basal segment naked. Their apical segment bears a strong spine on P₂-P₄. In addition, on P₃ only, a fine hair, slightly shorter than the spine, is seen. P₅: basoendopodites not fused, bearing three short, strong, blunt spines. Exopodites only slightly longer than wide, hardly prominent over the basoendopodites, and bearing two strong and two slim spines, subequal in length. The furcal rami are short, not longer than wide, tapering towards their tip, distinctly shorter than the anal segment. A row of spines stands at their ventral base. The anal operculum is not prominent, and bears five strong spines.

Relationship. — Geographically closest is *N. africana* Chappuis, 1955 from the beaches of Lake Tanganyika. Following PERKOVSKI's (1976) revision, it belongs to the *N. chappuisi*-group, which undoubtedly also


contains *N. somalica* on account of the presence of five appendages on the apical segment of the exopodite 3 of P₄. However, *N. africana* differs from *N. somalica* by its extremely long furcal rami, in addition to other structural details.



Figs. 2-10. — *Nitocrella somalica* n. sp. ♀: P₅ (Fig. 2), A₂ (Fig. 3), A₁ (Fig. 4), P₁ (Fig. 5), P₂ (Fig. 6), P₃ (Fig. 7), P₄ (Fig. 8), furcal rami (Fig. 9), furca and anal operculum (Fig. 10).

Morphologically closest appears to be *N. neutra* Kiefer, 1933, which is characterized by exactly the same spine and hair formula on the thoracopods 1-4, has short furcal rami, and a similar structure of P₅. Differences between both species are small and concern the relative length and size of the spines and hairs on P₅ and on the endopodite 2 of P₃. A very close relationship between both must thus be assumed, and without

broad geographical gap (possibly also Turkey) is not surprising.




Figs. 11-18. — *Metacyclops somalicus* n. sp. ♀: P₁ (Fig. 11), P₂ (Fig. 12), P₃ (Fig. 13), P₄ (Fig. 14), P₅ (Fig. 15), A₁ (Fig. 16), A₂ (Fig. 17), furca and anal operculum (Fig. 18).

Metacyclops somalicus n. sp.

Material examined. — Dissected female, mounted in alcohol, leg. SBS, 1.IX.1968.

Description. — Length 0.13 mm. Antenna 11-segmented. The spine formula 3.4.4.3. Probase of endopodite of P₁ 1. Apical segment of the endopodite 2 presenting a blunt tubercle.

ce of five appendages on the
 however, *N. africana* clearly
 g furcal rami, in addition to




10

Fig. 2), A₂ (Fig. 3), A₁ (Fig. 4), P₁
 furcal rami (Fig. 9), furca and anal
 10).

N. neutra Kiefer, 1933, which
 one and hair formula on the
 and a similar structure of P₅.
 and concern the relative length
 on the endopodite 2 of P₃. A
 thus be assumed, and without

the broad geographical gap (*N. neutra* is found in groundwaters of Yugo-
 slavia, possibly also Turkey), a specific distinction would have appeared
 doubtful.



Figs. 11-18. — *Metacyclops somalicus* n. sp. ♀: P₁ (Fig. 11), P₂ (Fig. 12),
 P₃ (Fig. 13), P₄ (Fig. 14), P₅ (Figs 15, 16), furca (Fig. 17), A₁ (Fig. 18).

SUBORDER CYCLOPOIDA

Metacyclops somalicus n. sp. (Figs 11-18).

Material examined. — Maas, well near the village; holotype (a fully
 dissected female, mounted in glycerin), 1 ♀ MF 012; paratypes 3 ♀♀ MF
 013; leg. SBS, 1.IX.1968.

Description. — Length 0.72 mm. A moderately small species. First
 antenna 11-segmented. The four thoracopods P₁-P₄ biarticulate and with
 spine formula 3.4.4.3. Precoxal lamella naked. Hair on basopodite at the
 base of endopodite of P₁ long and strongly built, reaching the tip of the
 apical segment of the endopodite. Inner angle of basopodite protruding,
 presenting a blunt tubercle between the inner angle and the insertion of

the endopodite. Endopodite 2 of P_4 with two apical spines; the shorter one is implanted outwardly and is less than half as long as the longer one, which is slightly sinuous and as long as the segment that bears it. P_5 as usual for the genus, with an isolated lateral seta and a single subcubic segment, bearing an apical seta and an apical spine. The latter is slightly longer than the segment which supports it. Shape of the receptaculum seminis not visible. Anal operculum rather conspicuous, rounded. Furcal rami about six times as long as wide with a very distinct notch at about 1/4 of the length from the base. Lateral seta inserted somewhat distal to the middle. Terminal setae of furca: inner one shorter than outer one, and outer one about as long as dorsal one. Of the long median setae, the inner one is distinctly longer than the outer one. Both are set with long and relatively spaced, stiff hairs.

Male unknown.

Relationship. — There is only one other species that presents a notch on the furcal rami, *M. tropicus* Kiefer, 1933. However, in this species the furcal rami are short, only about three times longer than wide.

Metacyclops chelazzii n. sp. (Figs 19-30).

Material examined. — Giohar (02°46'N-45°30'E), well in the garden of the SMOM (1); holotype (a fully dissected female, mounted in glycerin), 1 ♀ MF 014; paratypes, several ♀♀ MF 015, 1 ♂ MF 016; leg. SBS, 5.VIII.1968.


Extralimital material examined. — Rift valley Kenya, brooklet flowing into Lake Elmenteita, 1 ♀, leg. A. Coomans, 10.VIII.1975.

Derivatio nominis. — The species is named after Dr L. Chelazzi (Centro di Studio per la Faunistica ed Ecologia Tropicali del C.N.R., Florence), who collected it.

Descriptions. — *Female.* Length 1.1 mm. A moderately large species. First antenna 11-segmented. P_1 - P_4 biarticulate. Spine formula 3.4.4.3. Precoxal lamellae with two tubercles and, on P_2 - P_4 , with a transverse row of denticles. Hair at the base of endopodite 1 of P_1 not reaching beyond half the apical segment of the endopodite. Inner angle of basopodite rounded, but a strong thorn is found between the inner angle and the base of the endopodite. Exopodite 3 of P_4 with two apical spines; the inner one is longest and as long as the segment that bears it. The outer one is slightly over half as long as the inner one. P_5 consists of an isolated seta, and a segment which is as long as wide and bears two apical

(1) SMOM = Sovrano Militare Ordine di Malta.

appendages. The external seta, the isolated seta, and the lateral seta are only very slightly shorter than the segment that



Figs. 19-24. — *Metacyclops* sp. n.
P₄, second segment

supports it. Shape of the anal operculum well developed, rounded, tapering somewhat towards the base. Lateral seta implanted very distally, shorter than outer apical seta. Median seta longer than outer seta. Terminal setae strongly developed, rather closely approximated.

two apical spines; the shorter half as long as the longer one, the segment that bears it. P₅ as basal seta and a single subcubic apical spine. The latter is slightly shorter than the inner one. Shape of the receptaculum seminis conspicuous, rounded. Furcal rami very distinct notch at about 1/4 length inserted somewhat distal to the shorter than outer one, and the long median setae, the inner one. Both are set with long and

er species that presents a notch on P₃. However, in this species the furcal rami are longer than wide.

(N-45°30'E), well in the garden (collected female, mounted in glycerin), MF 015, 1 ♂ MF 016; leg. SBS,


t valley Kenya, brooklet flowing in the garden, 10.VIII.1975.

named after Dr L. Chelazzi (Centro di Studi e Ricerche in Zoologia e Botanica Tropicali del C.N.R., Florence),

mm. A moderately large species. Furcal rami crenulate. Spine formula 3.4.4.3. Furcal rami on P₂-P₄, with a transverse row of spinules. Inner angle of basopodite 1 of P₁ not reaching beyond middle. Inner angle of basopodite 1 of P₂-P₄ with two apical spines; the shorter one. P₅ consists of an isolated seta as wide and bears two apical

Malta.


appendages. The external one is a short seta, only about half as long as the isolated seta, and the internal one is a relatively strong spine, which is only very slightly shorter than the seta, but longer than the segment that



Figs. 19-24. — *Metacyclops chelazzii* n. sp. ♀: P₁ (Fig. 19), P₂ (Fig. 20), P₃ (Fig. 21), P₄, second specimen (Fig. 23), P₅, both sides (Fig. 24).

supports it. Shape of receptaculum seminis only partly visible. Anal operculum well developed, rounded. Furcal rami rather strongly built, tapering somewhat towards their apex, about 4.4 times as long as wide. Lateral seta implanted well distal to the middle. Dorsal seta distinctly shorter than outer apical seta, and about as long as inner apical seta. Inner median seta longer than outer median seta. Spinules on both long apical setae strongly developed, more than twice as long as the diameter of the setae, rather closely apposed.

Male. Length 0.9 mm. Differences with the female include the first antenna, which is 17-segmented and prehensile, and the structure of P₅, which shows an even more strongly developed apical spine on top of the



Figs. 25-30. — *Metacyclops chelazzii* n. sp.: P₅ and receptaculum seminis, ♀ (Fig. 25); P₅ and P₆, ♂ (Fig. 26); P₅, ♀ (Fig. 27); furca, ♀ (Fig. 28); A₁, ♂ (Fig. 29); A₁, ♀ (Fig. 30).

segment, and an apical seta of normal length (i.e. as long as the lateral seta). P₆ consists of a single spine.

Differential diagnosis. — The peculiar shape of P₅ will serve to differentiate this species from all its described congeners. Further, to the best of my knowledge, this is the first *Metacyclops* with a male in which P₆ is reduced to a single appendix, all other species in which the male is known showing either two or three appendages on P₆.

Metacyclops sp.

Material examined. — Baidoa, well of the aqueduct, several copepodites MF 018, leg. SBS, 27.VIII.1968.

Afrocylops gibsoni (Brady)

Material examined. — MF 017, leg. SBS, 5.VIII.

Remarks. — A wide range of climatic zones of the continent logically be expected to occur.

Two species of Cladocera among which are new to science and *Metacyclops chelazzii*, geographical relationships are


L'Autore ha studiato un'e quattro specie di Copepodi: *Nitocrella somalica*, *Metacyclops chelazzii*, e due altre sconosciute provenienti dalla Somalia. Sono descritte con disegni grafiche.

- CHAPPUIS, P. A. 1955. Harpacticoides. Bot. afr. 51: 68-80.
 DADAY, E. VON 1907. Plancton der Bahr- und Sobataneen. Zool. Anz. 25: 245-262.
 DUMONT, H. J. 1979. Limnology of the Baidoa well. Limnol. Oceanogr. 24: 100-104.
 KIEFER, F. 1933a. Voyage de C. G. S. en Somalie française. Freilebende Binnentiere. Arch. Hydrobiol. 28: 121-130.
 KIEFER, F. 1933b. Neue Süßwasserharpacticoiden. Zool. Anz. 309-318.
 PARDI, L. 1976. L'attività del Copepode *Nitocrella somalica* (Chapuis) (Copepoda, Harpacticoida) in «Tropicali» del Consiglio Nazionale delle Ricerche. Monitore zool. ital. (N. S.) 10: 1-10.
 PETKOVSKI, T. K. 1976. Drei neue Copepodenarten des Genus *Nitocrella* Chapuis (Copepoda, Harpacticoida). Monitore zool. ital. (N. S.) 10: 126: 26 pp.

Address of the author: Dr. HENRI DUMONT, Ledeganckstraat 35, Gent (Belgium).

Monitore zool. ital. (N. S.) Suppl.

the female include the first
le, and the structure of P₅,
d apical spine on top of the



receptaculum seminis, ♀ (Fig. 25);
(Fig. 28); A₁, ♂ (Fig. 29); A₁,

i.e. as long as the lateral seta).

r shape of P₅ will serve to
ed congeners. Further, to the
cyclops with a male in which
species in which the male is
ges on P₆.

of the aqueduct, several co-

Afrocyclops gibsoni (Brady, 1904).

Material examined. — Giohar, well in the garden of the SMOM, 1 ♀
MF 017, leg. SBS, 5.VIII.1968.

Remarks. — A widespread African species, known from virtually all
climatic zones of the continent, including the Sahara. It could therefore
logically be expected to occur in Somalia as well.

SUMMARY

Two species of Cladocera and four species of Copepoda — three
among which are new to science (*Nitocrella somalica*, *Metacyclops somalicus*
and *Metacyclops chelazzii*) — are reported from Somalia. Their bio-
geographical relationships are briefly discussed.

RIASSUNTO

L'Autore ha studiato una piccola collezione di due specie di Cladocera
e quattro specie di Copepoda (tre delle quali sono nuove per la scienza:
Nitocrella somalica, *Metacyclops somalicus* e *Metacyclops chelazzii*) prove-
nienti dalla Somalia. Sono brevemente discusse le loro relazioni biogeo-
grafiche.

REFERENCES

- CHAPPUIS, P. A. 1955. Harpacticoides psammiques du lac Tanganika. *Revue Zool. Bot. afr.* 51: 68-80.
- DADAY, E. VON 1907. Plancton-Tiere aus dem Victoria Nuanza. *Zool. Jb. (Syst.)* 25: 245-262.
- DUMONT, H. J. 1979. *Limnologie van Sahara en Sahel*. University of Ghent, 557 pp.
- KIEFER, F. 1933a. Voyage de Ch. Alluaud et P. A. Chappuis en Afrique occidentale française. Freilebende Binnengewässer Copepoden. Diaptomiden und Cyclopiden. *Arch. Hydrobiol.* 28: 121-142.
- KIEFER, F. 1933b. Neue Süßwasser Copepoden aus Jugoslawien. *Zool. Anz.* 101: 309-318.
- PARDI, L. 1976. L'attività del « Centro di Studio per la Faunistica ed Ecologia Tropicali » del Consiglio Nazionale delle Ricerche nel quinquennio 1971-1976. *Monitore zool. ital. (N. S.) Suppl.* 7: 195-269.
- PETKOVSKI, T. K. 1976. Drei neue *Nitocrella*-Arten aus Kuba, zugleich eine Revision des Genus *Nitocrella* Chappuis (s. restr.). *Acta Mus. maced. Sci. nat.* 15 (1) 126: 26 pp.

Address of the author: Dr. HENRI J. DUMONT, Zoological Institute, University of Gent, Ledeganckstraat 35, Gent (Belgium).

Monitore zool. ital. (N. S.) Suppl. XIV: 103-111, 1981